

Train faster, generalize better: Stability of stochastic gradient descent

Moritz Hardt*

Benjamin Recht[†]

Yoram Singer[‡]

September 4, 2015

Abstract

We show that any model trained by a stochastic gradient method with few iterations has vanishing generalization error. We prove this by showing the method is algorithmically stable in the sense of Bousquet and Elisseeff. Our analysis only employs elementary tools from convex and continuous optimization. Our results apply to both convex and non-convex optimization under standard Lipschitz and smoothness assumptions.

Applying our results to the convex case, we provide new explanations for why multiple epochs of stochastic gradient descent generalize well in practice. In the nonconvex case, we provide a new interpretation of common practices in neural networks, and provide a formal rationale for stability-promoting mechanisms in training large, deep models. Conceptually, our findings underscore the importance of reducing training time beyond its obvious benefit.

1 Introduction

The most widely used optimization method in machine learning practice is *stochastic gradient method* (SGM). Stochastic gradient methods aim to minimize the empirical risk of a model by repeatedly computing the gradient of a loss function on a single training example, or a batch of few examples, and updating the model parameters accordingly. SGM is scalable, robust, and performs well across many different domains ranging from smooth and strongly convex problems to complex non-convex objectives.

In a nutshell, our results establish that:

Any model trained with stochastic gradient method in a reasonable amount of time attains small generalization error.

As training time is inevitably limited in practice, our results help to explain the strong generalization performance of stochastic gradient methods observed in practice. More concretely, we bound the generalization error of a model in terms of the number of iterations that stochastic gradient method took in order to train the model. Our main analysis tool is to employ the notion of algorithmic stability due to Bousquet and Elisseeff [4]. We demonstrate that the stochastic gradient method is

*Email: mrtz@google.com

[†]Email: brecht@berkeley.edu, work performed at Google.

[‡]Email: singer@google.com

stable provided that the objective is relatively smooth and the number of steps taken is sufficiently small.

It is common in practice to perform a linear number of steps in the size of the sample and to access each data point multiple times. Our results show in a broad range of settings that, provided the number of iterations is linear in the number of data points, the generalization error is bounded by a vanishing function of the sample size. The results hold true even for complex models with large number of parameters and no explicit regularization term in the objective. Namely, fast training time by itself is sufficient to prevent overfitting.

Our bounds are algorithm specific: Since the number of iterations we allow can be larger than the sample size, an arbitrary algorithm could easily achieve small training error by memorizing all training data with no generalization ability whatsoever. In contrast, if the stochastic gradient method manages to overfit the training data in a reasonable number of iterations, it is still guaranteed to generalize.

Conceptually, we show that minimizing training time is not only beneficial for the obvious computational advantages, but in addition has the important by-product of decreasing generalization error. Consequently, it may make sense for practitioners to focus on minimizing training time, for instance, by designing model architectures for which stochastic gradient method converges fastest to a desired error level.

1.1 Our contributions

Our focus is on generating *generalization bounds* for models learned with stochastic gradient descent. Recall that the generalization bound is the expected difference between the error a model incurs on a training set versus the error incurred on a new data point, sampled from the same distribution that generated the training data. Throughout, we assume we are training models using n sampled data points.

Our results build on a fundamental connection between the generalization error of an algorithm and its *stability* properties. Roughly speaking, an algorithm is *stable* if the training error it achieves varies only slightly if we change any single training data point. The precise notion of stability we use is known as *uniform stability* due to [4]. It states that a randomized algorithm A is uniformly stable if for all data sets differing in only one element, the learned models produce nearly the same predictions. We review this method in Section 2, and provide a new adaptation of this theory to *iterative* algorithms.

In Section 3, we show that stochastic gradient is uniformly stable, and our techniques mimic its convergence proofs. For convex loss functions, we prove that the stability measure decreases as a function of the sum of the step sizes. For strongly convex loss functions, we show that stochastic gradient is stable, even if we train for an arbitrarily long time.

More surprisingly, our results carry over to the case where the loss-function is non-convex. In this case we show that the method generalizes provided the steps are sufficiently small and the number of iterations is not too large. More specifically, we show the number of steps of stochastic gradient can grow as n^c for a small $c > 1$. This provides some explanation as to why neural networks can be trained for multiple epochs of stochastic gradient and still exhibit excellent generalization. In Section 4, we furthermore show that various heuristics used in practice, especially in the deep learning community, help to increase the stability of stochastic gradient method. For example, the popular dropout scheme [19, 39] improves all of our bounds. Similarly, ℓ_2 -regularization improves

the exponent of n in our non-convex result. In fact, we can drive the exponent arbitrarily close to $1/2$ while preserving the non-convexity of the problem.

We can combine our bounds on the generalization error of stochastic gradient method with optimization bounds quantifying the convergence of the empirical loss achieved by SGM. In Section 5, we show that models trained for multiple epochs match classic bounds for stochastic gradient [27,28].

1.2 Related work

There is a venerable line of work on stability and generalization dating back more than thirty years [4, 8, 18, 25, 38]. The landmark work by Bousquet and Elisseeff [4] introduced the notion of *uniform stability* that we rely on. They showed that several important classification techniques are uniformly stable. In particular, under certain regularity assumptions, it was shown that the *optimizer* of a regularized empirical loss minimization problem is uniformly stable. Previous work generally applies only to the exact minimizer of specific optimization problems. It is not immediately evident on how to compute a generalization bound for an approximate minimizer such as one found by using stochastic gradient. Subsequent work studied stability bounds for randomized algorithms but focused on random perturbations of the cost function, such as those induced by bootstrapping or bagging [9]. This manuscript differs from this foundational work in that it derives stability bounds about the learning *procedure*, analyzing algorithmic properties that induce stability.

Stochastic gradient descent, of course, is closely related to our inquiry. Classic results by Nemirovski and Yudin show that the stochastic gradient method produces are nearly optimal for empirical risk minimization of convex loss functions [11,26–28]. These results have been extended by many machine learning researchers, yielding tighter bounds and probabilistic guarantees [13,14,34]. However, there is an important limitation of all of this prior art. The derived generalization bounds only hold for single passes over the data. That is, in order for the bounds to be valid, each training example must be used no more than once in a stochastic gradient update. In practice, of course, one tends to run multiple *epochs* of the stochastic gradient method. Our results resolve this issue by combining stability with optimization error. We use the foundational results to estimate the error on the *empirical risk* and then use stability to derive a deviation from the true risk. This enables us to study the risk incurred by multiple epochs and provide simple analyses of regularization methods for convex stochastic gradient. We compare our results to this related work in Section 5. We note that Rosasco and Villa obtain risk bounds for least squares minimization with an incremental gradient method in terms of the number of epochs [36]. These bounds are akin to our study in Section 5, although our results are incomparable due to various different assumptions.

Finally, we note that in the non-convex case, the stochastic gradient method is remarkably successful for training large neural networks [2, 19]. However, our theoretical understanding of this method is limited. Several authors have shown that the stochastic gradient method finds a stationary point of nonconvex cost functions [12,21]. Beyond asymptotic convergence to stationary points, little is known about finding models with low training or generalization error in the nonconvex case. There have recently been several important studies investigating optimal training of neural nets. For example Livni *et al.* show that networks with polynomial activations can be learned in a greedy fashion [24]. Janzamin *et al.* [16] show that two layer neural networks can be learned using tensor methods. Arora *et al.* [1] show that two-layer sparse coding dictionaries can be learned via stochastic gradient. Our work complements these developments: rather than providing new insights into mechanisms that yield low training error, we provide insights into mechanisms

that yield low generalization error. If one can achieve low training error quickly on a nonconvex problem with stochastic gradient, our results guarantee that the resulting model generalizes well.

2 Stability of randomized iterative algorithms

Consider the following general setting of supervised learning. There is an unknown joint distribution \mathcal{D} over pairs of labeled examples from a product space $X \times Y$. We receive a sample $S = (z_1, \dots, z_n)$ of n examples drawn i.i.d. from \mathcal{D} . Our goal is to find a model w with small *population risk*, defined as:

$$R[w] \stackrel{\text{def}}{=} \mathbb{E}_{z \sim \mathcal{D}} f(w; z).$$

Here, where f is a *loss function* and $f(w; z)$ designates the *loss* of the model described by w encountered on example z .

Since we cannot measure the objective $R[w]$ directly, we instead use a sample-averaged proxy, the *empirical risk*, defined as

$$R_S[w] \stackrel{\text{def}}{=} \frac{1}{n} \sum_{i=1}^n f(w; z_i),$$

The *generalization error* of a model w is the difference $R_S[w] - R[w]$. When $w = A(S)$ is chosen as a function of the data by a potentially randomized algorithm A it makes sense to consider the expected generalization error

$$\epsilon_{\text{gen}} \stackrel{\text{def}}{=} \mathbb{E}[R_S[A(S)] - R[A(S)]], \quad (2.1)$$

where the expectation is over the randomness of A and the sample S .

In order to bound the generalization error of an algorithm, we employ with the following notion of *uniform stability* in which we allow randomized algorithms as well.

Definition 2.1. *A randomized algorithm A is ϵ -uniformly stable if for all data sets $S, S' \in Z^n$ such that S and S' differ in at most one example, we have*

$$\sup_z \mathbb{E}_A [f(A(S); z) - f(A(S'); z)] \leq \epsilon. \quad (2.2)$$

Here, the expectation is taken only over the internal randomness of A . We will denote by $\epsilon_{\text{stab}}(A, n)$ the infimum over all ϵ for which (2.2) holds. We will omit the tuple (A, n) when it is clear from the context.

We recall the important theorem that uniform stability implies *generalization in expectation*. Since our notion of stability differs slightly from existing ones with respect to the randomness of the algorithm, we include a proof for the sake of completeness. The proof is very similar to the argument in Lemma 7 of [4].

Theorem 2.2 (Generalization in expectation). *Let A be ϵ -uniformly stable. Then,*

$$|\mathbb{E}_{S,A} [R_S[A(S)] - R[A(S)]]| \leq \epsilon.$$

Proof. Denote by $S = (z_1, \dots, z_n)$ and $S' = (z'_1, \dots, z'_n)$ two independent random samples and let $S^{(i)} = (z_1, \dots, z_{i-1}, z'_i, z_{i+1}, \dots, z_n)$ be the sample that is identical to S except in the i 'th example where we replace z_i with z'_i . With this notation, we get that

$$\begin{aligned} \mathbb{E}_S \mathbb{E}_A [R_S[A(S)]] &= \mathbb{E}_S \mathbb{E}_A \left[\frac{1}{n} \sum_{i=1}^n f(A(S); z_i) \right] \\ &= \mathbb{E}_S \mathbb{E}_{S'} \mathbb{E}_A \left[\frac{1}{n} \sum_{i=1}^n f(A(S^{(i)}); z'_i) \right] \\ &= \mathbb{E}_S \mathbb{E}_{S'} \mathbb{E}_A \left[\frac{1}{n} \sum_{i=1}^n f(A(S); z'_i) \right] + \delta \\ &= \mathbb{E}_S \mathbb{E}_A [R[A(S)]] + \delta, \end{aligned}$$

where we can express δ as

$$\delta = \mathbb{E}_S \mathbb{E}_{S'} \mathbb{E}_A \left[\frac{1}{n} \sum_{i=1}^n f(A(S^{(i)}); z'_i) - \frac{1}{n} \sum_{i=1}^n f(A(S); z'_i) \right].$$

Furthermore, taking the supremum over any two data sets S, S' differing in only one sample, we can bound the difference as

$$|\delta| \leq \sup_{S, S', z} \mathbb{E}_A [f(A(S); z) - \mathbb{E}_A f(A(S'); z)] \leq \epsilon,$$

by our assumption on the uniform stability of A . The claim follows. \square

Theorem 2.2 proves that if an algorithm is uniformly stable, then its generalization is small. We now turn to some properties of iterative algorithms that control their uniform stability.

2.1 Properties of update rules

We consider general update rules of the form $G: \Omega \rightarrow \Omega$ which map a point $w \in \Omega$ in the parameter space to another point $G(w)$. The most common update is the gradient update rule

$$G(w) = w - \alpha \nabla f(w),$$

where $\alpha \geq 0$ is a step size and $f: \Omega \rightarrow \mathbb{R}$ is a function that we want to optimize.

The canonical update rule we will consider in this manuscript is an incremental gradient update, where $G(w) = w - \alpha \nabla f(w)$ for some convex function f . We will return to a detailed discussion of this specific update in the sequel, but the reader should keep this particular example in mind throughout the remainder of this section.

The following two definitions provide the foundation of our analysis of how two different sequences of update rules diverge when iterated from the same starting point. These definitions will ultimately be useful when analyzing the stability of stochastic gradient descent.

Definition 2.3. *An update rule is η -expansive if*

$$\sup_{v, w \in \Omega} \frac{\|G(v) - G(w)\|}{\|v - w\|} \leq \eta. \tag{2.3}$$

Definition 2.4. An update rule is σ -bounded if

$$\sup_{w \in \Omega} \|w - G(w)\| \leq \sigma. \quad (2.4)$$

With these two properties, we can establish the following lemma of how a sequence of updates to a model diverge when the training set is perturbed.

Lemma 2.5 (Growth recursion). *Fix an arbitrary sequence of updates G_1, \dots, G_T and another sequence G'_1, \dots, G'_T . Let $w_0 = w'_0$ be a starting point in Ω and define $\delta_t = \|w'_t - w_t\|$ where w_t, w'_t are defined recursively through*

$$w_{t+1} = G_t(w_t) \quad w'_{t+1} = G_t(w'_t). \quad (t > 0)$$

Then, we have the recurrence relation

$$\begin{aligned} \delta_0 &= 0 \\ \delta_{t+1} &\leq \begin{cases} \eta\delta_t & G_t = G'_t \text{ is } \eta\text{-expansive} \\ \min(\eta, 1)\delta_t + 2\sigma_t & G_t \text{ and } G'_t \text{ are } \sigma\text{-bounded, } G_t \text{ is } \eta \text{ expansive} \end{cases} \quad (t > 0) \end{aligned}$$

Proof. The first bound on δ_t follow directly from the assumption that $G_t = G'_t$ and the definition of expansiveness. For the second bound, recall from Definition 2.4 that if G_t and G'_t are σ -bounded, then by the triangle inequality,

$$\begin{aligned} \delta_{t+1} &= \|G(w_t) - G'(w'_t)\| \\ &\leq \|G(w_t) - w_t + w_t - w'_t + G'(w'_t)\| + \|w_t - w'_t\| \\ &\leq \delta_t + \|G(w_t) - w_t\| + \|G'(w'_t) - w'_t\| \\ &\leq \delta_t + 2\sigma, \end{aligned}$$

which gives half of the second bound. We can alternatively bound δ_{t+1} as

$$\begin{aligned} \delta_{t+1} &= \|G_t(w_t) - G'_t(w'_t)\| \\ &= \|G_t(w_t) - G_t(w'_t) + G_t(w'_t) - G'_t(w'_t)\| \\ &\leq \|G_t(w_t) - G_t(w'_t)\| + \|G_t(w'_t) - G'_t(w'_t)\| \\ &\leq \|G_t(w_t) - G_t(w'_t)\| + \|w'_t - G_t(w'_t)\| + \|w'_t - G'_t(w'_t)\| \\ &\leq \eta\delta_t + 2\sigma. \end{aligned}$$

□

We now turn to using these lemmas to analyze one of the core algorithms in statistical learning: the stochastic gradient method.

3 Stability of Stochastic Gradient Descent

Given n labeled examples $S = (z_1, \dots, z_n)$ where $z_i = (x_i, y_i) \in X \times Y$, consider a *decomposable* objective function

$$f(w) = \frac{1}{n} \sum_{i=1}^n f(w; (x_i, y_i)),$$

where $f(w; (x_i, y_i))$ denotes the *loss* of w on the example (x_i, y_i) . The stochastic gradient update for this problem with *learning rate* $\alpha_t > 0$ is given by

$$w_{t+1} = w_t - \alpha_t \nabla_w f(w_t; z_{i_t}).$$

Stochastic gradient method (SGM) is the algorithm resulting from performing stochastic gradient updates T times where the indices i_t are randomly chosen. There are two popular schemes for choosing the examples' indices. One is to pick i_t uniformly at random in $\{1, \dots, n\}$ at each step. The other is to choose a random permutation over $\{1, \dots, n\}$ and cycle through the examples repeatedly in the order determined by the permutation. Our results hold for both variants.

In parallel with the previous section the stochastic gradient method is akin to applying the *gradient update rule* defined as follows.

Definition 3.1. For a nonnegative step size $\alpha \geq 0$ and a function $f: \Omega \rightarrow \mathbb{R}$, we define the gradient update rule $G_{f,\alpha}$ as

$$G_{f,\alpha}(w) = w - \alpha \nabla f(w).$$

3.1 Proof idea: Stability of stochastic gradient method

In order to prove that the stochastic gradient method is stable, we will analyze the output of the algorithm on two data sets that differ in precisely one location. Note that if the loss function is L -Lipschitz, we have $\mathbb{E} |f(w; \cdot) - f(w'; \cdot)| \leq L \mathbb{E} \|w - w'\|$ for any w and w' . Hence, it suffices to analyze how w_t and w'_t diverge in the domain as a function of time t . Recalling that w_t is obtained from w_{t-1} via a gradient update, our goal is to bound $\delta_t = \|w_t - w'_t\|$ recursively and in expectation as a function of δ_{t-1} .

There are two cases to consider. In the first case, SGM selects the index of an example at step t on which is identical in S and S' . In this case, the gradients added to w_t and w'_t are identical. Unfortunately, it could still be the case that δ_t grows. Below, we will show how to control δ_t in terms of the convexity and smoothness properties of the stochastic gradients.

The second case to consider is when SGM selects the one example to update in which S and S' differ. Note that this happens only with probability $1/n$ if examples are selected randomly. In this case, we simply bound the increase in δ_t by the norm of the two gradient $\nabla f(w_{t-1}, z)$ and $\nabla f(w'_{t-1}, z')$. The sum of the norms is bounded by $2\alpha_t L$ and we obtain $\delta_t \leq \delta_{t-1} + 2\alpha_t L$. Combining the two cases, we can then solve a simple recurrence relation to obtain a bound on δ_T .

This simple approach suffices to obtain the desired result in the convex case, but there are additional difficulties in the non-convex case. Here, we need to use an intriguing stability property of stochastic gradient method. Specifically, the first time step t_0 at which SGM even encounters the example in which S and S' differ is a random variable in $\{1, \dots, n\}$ which tends to be relatively large. Specifically, for any $m \in \{1, \dots, n\}$, the probability that $t_0 \leq m$ is upper bounded by m/n . This allows us to argue that SGM has a long “burn-in period” where δ_t does not grow at all. Once δ_t begins to grow, the step size has already decayed allowing us to obtain a non-trivial bound.

We now turn to making this argument precise.

3.2 Expansion properties of stochastic gradients

Let us now record some of the core properties of the stochastic gradient update. The gradient update rule is bounded provided that the function f satisfies the following common Lipschitz condition.

Definition 3.2. We say that f is L -Lipschitz if for all points u in the domain of f we have $\|\nabla f(x)\| \leq L$. This implies that

$$|f(u) - f(v)| \leq L\|u - v\|. \quad (3.1)$$

Lemma 3.3. Assume that f is L -Lipschitz. Then, the gradient update $G_{f,\alpha}$ is (αL) -bounded.

Proof. By our Lipschitz assumption, $\|w - G_{f,\alpha}(w)\| = \|\alpha \nabla f(w)\| \leq \alpha L$. \square

We now turn to expansiveness. As we will see shortly, different expansion properties are achieved for non-convex, convex, and strongly convex functions.

Definition 3.4. A function $f: \Omega \rightarrow \mathbb{R}$ is convex if for all $u, v \in \Omega$ we have

$$f(u) \geq f(v) + \langle \nabla f(v), u - v \rangle.$$

Definition 3.5. A function $f: \Omega \rightarrow \mathbb{R}$ is γ -strongly convex if for all $u, v \in \Omega$ we have

$$f(u) \geq f(v) + \langle \nabla f(v), u - v \rangle + \frac{\gamma}{2}\|u - v\|^2.$$

The following standard notion of smoothness leads to a bound on how expansive the gradient update is.

Definition 3.6. A function $f: \Omega \rightarrow \mathbb{R}$ is β -smooth if for all $u, v \in \Omega$ we have

$$\|\nabla f(u) - \nabla f(v)\| \leq \beta\|u - v\|. \quad (3.2)$$

In general, smoothness will imply that the gradient updates cannot be overly expansive. When the function is also convex and the step size is sufficiently small the gradient update becomes non-expansive. When the function is additionally strongly convex, the gradient update becomes *contractive* in the sense that η will be less than one and u and v will actually shrink closer to one another. The majority of the following results can be found in several textbooks and monographs. Notable references are Polyak [33] and Nesterov [29]. We include proofs in the appendix for completeness.

Lemma 3.7. Assume that f is β -smooth. Then, the following properties hold.

1. $G_{f,\alpha}$ is $(1 + \alpha\beta)$ -expansive.
2. Assume in addition that f is convex. Then, for any $\alpha \leq 2/\beta$, the gradient update $G_{f,\alpha}$ is 1-expansive.
3. Assume in addition that f is γ -strongly convex. Then, for $\alpha \leq \frac{2}{\beta+\gamma}$, $G_{f,\alpha}$ is $\left(1 - \frac{2\alpha\beta\gamma}{\beta+\gamma}\right)$ -expansive.

We will also make use of the following convenient lemma for analyzing the stability of the stochastic gradient method.

Lemma 3.8. *Assume that the loss function $f(\cdot; z)$ is L -Lipschitz for all z . Let S and S' be two samples of size n differing in only a single example. Denote by w_T and w'_T the output of T steps of SGM on S and S' , respectively. Then, for every $z \in X \times Y$ and every $t_0 \in \{0, 1, \dots, n\}$, under both the random update rule and the random permutation rule, we have*

$$\mathbb{E} |f(w_T; z) - f(w'_T; z)| \leq \frac{t_0}{n} + L \mathbb{E} [\delta_T \mid \delta_{t_0} = 0] .$$

Proof. Let S and S' be two samples of size n differing in only a single example, and let $z \in X \times Y$ be an arbitrary example. Consider running SGM on sample S and S' , respectively. As stated, w_T and w'_T denote the corresponding outputs of SGM. Let $\mathcal{E} = \mathbf{1}[\delta_{t_0} = 0]$ denote the event that $\delta_{t_0} = 0$. We have,

$$\begin{aligned} \mathbb{E} |f(w_T; z) - f(w'_T; z)| &= \mathbb{P} \{ \mathcal{E} \} \mathbb{E} [|f(w_T; z) - f(w'_T; z)| \mid \mathcal{E}] \\ &\quad + \mathbb{P} \{ \mathcal{E}^c \} \mathbb{E} [|f(w_T; z) - f(w'_T; z)| \mid \mathcal{E}^c] \\ &\leq \mathbb{E} [|f(w_T; z) - f(w'_T; z)| \mid \mathcal{E}] + \mathbb{P} \{ \mathcal{E}^c \} \\ &\leq L \mathbb{E} [\|w_T - w'_T\| \mid \mathcal{E}] + \mathbb{P} \{ \mathcal{E}^c \} . \end{aligned}$$

The first inequality uses the fact that $|f(w_T; z) - f(w'_T; z)| \leq 1$, from our assumption on the range of the function f . The second inequality follows from the Lipschitz assumption.

It remains to bound $\mathbb{P} \{ \mathcal{E}^c \}$. Toward that end, let $i^* \in \{1, \dots, n\}$ denote the position in which S and S' differ and consider the random variable I assuming the index of the first time step in which SGM uses the example z_{i^*} . Note that when $I > t_0$, then we must have that $\delta_{t_0} = 0$, since the execution on S and S' is identical until step t_0 . Hence,

$$\mathbb{P} \{ \mathcal{E}^c \} = \mathbb{P} \{ \delta_{t_0} \neq 0 \} \leq \mathbb{P} \{ I \leq t_0 \} .$$

Under the random permutation rule, I is a uniformly random number in $\{1, \dots, n\}$ and therefore

$$\mathbb{P} \{ I \leq t_0 \} = \frac{t_0}{n} .$$

This proves the claim we stated for the random permutation rule. For the random selection rule, we have by the union bound $\mathbb{P} \{ I \leq t_0 \} \leq \sum_{t=1}^{t_0} \mathbb{P} \{ I = t \} = \frac{t_0}{n}$. This completes the proof. \square

Henceforth we will no longer mention which random selection rule we use as the proofs are almost identical for both rules.

3.3 Convex optimization

We begin with a simple stability bound for convex loss minimization via stochastic gradient method.

Theorem 3.9. *Assume that the loss function $f(\cdot; z) \in [0, 1]$ is β -smooth, convex and L -Lipschitz for every z . Suppose that we run SGM with step sizes $\alpha_t \leq 2/\beta$ for T steps. Then, SGM satisfies uniform stability with*

$$\epsilon_{\text{stab}} \leq \frac{2L^2}{n} \sum_{t=1}^T \alpha_t .$$

Proof. Let S and S' be two samples of size n differing in only a single example. Consider the gradient updates G_1, \dots, G_T and G'_1, \dots, G'_T induced by running SGM on sample S and S' , respectively. Let w_T and w'_T denote the corresponding outputs of SGM.

We now fix an example $z \in X \times Y$ and apply Lemma 3.8 to $f(\cdot; z)$ with $t_0 = 0$, to get

$$\mathbb{E} |f(w_T; z) - f(w'_T; z)| \leq L \mathbb{E} [\delta_T | \delta_0 = 0] = L \mathbb{E} [\delta_T], \quad (3.3)$$

where $\delta_t = \|w_T - w'_T\|$. Observe that at step t , with probability $1 - 1/n$, the example selected by SGM is the same in both S and S' . In this case we have that $G_t = G'_t$ and we can use the 1-expansivity of the update rule G_t which follows from Lemma 3.7.2 using the fact that the objective function is convex and that $\alpha_t \leq 2/\beta$. With probability $1/n$ the selected example is different in which case we use that both G_t and G'_t are $\alpha_t L$ -bounded as a consequence of Lemma 3.3. Hence, we can apply Lemma 2.5 and linearity of expectation to conclude that for every t ,

$$\mathbb{E} [\delta_{t+1}] \leq \left(1 - \frac{1}{n}\right) \mathbb{E} [\delta_t] + \frac{1}{n} \mathbb{E} [\delta_t] + \frac{2\alpha_t L}{n} = \mathbb{E} [\delta_t] + \frac{2L\alpha_t}{n}. \quad (3.4)$$

Unraveling the recursion gives

$$\mathbb{E} [\delta_T] \leq \frac{2L}{n} \sum_{t=1}^T \alpha_t.$$

Plugging this back into equation (3.3), we obtain

$$\mathbb{E} |f(w_T; z) - f(w'_T; z)| \leq \frac{2L^2}{n} \sum_{t=1}^T \alpha_t.$$

Since this bound holds for all S, S' and z , we obtain the desired bound on the uniform stability. \square

3.4 Strongly convex optimization

In the strongly convex case our bound has no dependence on the number of steps at all.

Theorem 3.10. *Assume that the loss function $f(\cdot; z) \in [0, 1]$ is γ -strongly convex, L -Lipschitz and β -smooth for all z . Suppose we run SGM with constant step size $\alpha \leq 1/\beta$ for T steps. Then, SGM satisfies uniform stability with*

$$\epsilon_{\text{stab}} \leq \frac{2L^2}{\gamma n}.$$

Proof. The proof is analogous to that of Theorem 3.9 with a slightly different recurrence relation. We repeat the the argument of completeness. Let S and S' be two samples of size n differing in only a single example. Consider the gradient updates G_1, \dots, G_T and G'_1, \dots, G'_T induced by running SGM on sample S and S' , respectively. Let w_T and w'_T denote the corresponding outputs of SGM.

Applying Lemma 3.8 with $t_0 = 0$, we obtain

$$\mathbb{E} |f(w_T; z) - f(w'_T; z)| \leq L \mathbb{E} [\delta_T | \delta_0 = 0] = L \mathbb{E} [\delta_T], \quad (3.5)$$

where $\delta_t = \|w_T - w'_T\|$. Observe that at step t , with probability $1 - 1/n$, the example selected by SGM is the same in both S and S' . In this case we have that $G_t = G'_t$. We can now apply the following useful simplification of Lemma 3.7.3 if $\alpha \leq 1/\beta$. Since we also require that $\alpha \leq \frac{2}{\beta+\gamma}$ we

get that $\frac{2\alpha\beta\gamma}{\beta+\gamma} \geq \alpha^2\beta\gamma \geq \alpha\gamma$, thus $G_{f,\alpha}$ is $(1 - \alpha\gamma)$ -expansive. With probability $1/n$ the selected example is different in which case we use that both G_t and G'_t are αL -bounded as a consequence of Lemma 3.3 which appears in the next section. Hence, we can apply Lemma 2.5 and linearity of expectation to conclude that for every t ,

$$\begin{aligned}\mathbb{E} \delta_{t+1} &\leq \left(1 - \frac{1}{n}\right) (1 - \alpha\gamma) \mathbb{E} \delta_t + \frac{1}{n}(1 - \alpha\gamma) \mathbb{E} \delta_t + \frac{2\alpha L}{n} \\ &= (1 - \alpha\gamma) \mathbb{E} \delta_t + \frac{2\alpha L}{n}\end{aligned}\tag{3.6}$$

Unraveling the recursion gives

$$\mathbb{E} \delta_T \leq \frac{2L\alpha}{n} \sum_{t=0}^T (1 - \alpha\gamma)^t \leq \frac{2L}{\gamma n}.$$

Plugging the above inequality into equation (3.3), we obtain

$$\mathbb{E} |f(w_T; z) - f(w'_T; z)| \leq \frac{2L^2}{\gamma n}.$$

Since this bounds holds for all S, S' and z , the lemma follows. \square

We would like to note that a nearly identical result holds for a “staircase” decaying step-size that is also popular in machine learning and stochastic optimization.

Theorem 3.11. *Assume that the loss function $f(\cdot; z) \in [0, 1]$ is γ -strongly convex, L -Lipschitz, and β -smooth function for all z . Suppose we run SGM with step sizes $\alpha_t = \frac{1}{\gamma t}$. Then, SGM has uniform stability of*

$$\epsilon_{\text{stab}} \leq \frac{2L^2 + \beta}{\gamma n}.$$

Proof. Note that once $t > \frac{\beta}{\gamma}$, the iterates are contractive with contractivity $1 - \alpha_t\gamma \leq 1 - \frac{1}{t}$. Thus, for $t \geq t_0 := \frac{\beta}{\gamma}$, we have

$$\begin{aligned}\mathbb{E}[\delta_{t+1}] &\leq \left(1 - \frac{1}{n}\right)(1 - \alpha_t\gamma) \mathbb{E}[\delta_t] + \frac{1}{n}((1 - \alpha_t\gamma) \mathbb{E}[\delta_t] + 2\alpha_t L) \\ &= (1 - \alpha_t\gamma) \mathbb{E}[\delta_t] + \frac{2\alpha_t L}{n} \\ &= \left(1 - \frac{1}{t}\right) \mathbb{E}[\delta_t] + \frac{2L}{\gamma t n}.\end{aligned}$$

Assuming that $\delta_{t_0} = 0$ and expanding this recursion, we find.

$$\mathbb{E}[\delta_T] \leq \sum_{t=t_0}^T \left\{ \prod_{s=t+1}^T \left(1 - \frac{1}{s}\right) \right\} \frac{2L}{\gamma t n} = \sum_{t=t_0}^T \frac{t}{T} \frac{2L}{\gamma t n} = \frac{T - t_0}{T} \frac{2L}{\gamma n}.$$

Now, the result follows from Lemma 3.8 while the fact that $t_0 = \frac{\beta}{\gamma}$. \square

3.5 Non-convex optimization

In this section we prove stability results for stochastic gradient methods that do not require convexity. We will still assume that the objective function is smooth and Lipschitz as defined previously.

Theorem 3.12. *Assume that $f(\cdot; z) \in [0, 1]$ is an L -Lipschitz and β -smooth loss function for every z . Suppose that we run SGM for T steps with monotonically non-increasing step sizes $\alpha_t \leq c/t$. Then, SGM has uniform stability with*

$$\epsilon_{\text{stab}} \leq \frac{1 + 1/\beta c}{n - 1} (2cL^2)^{\frac{1}{\beta c + 1}} T^{\frac{\beta c}{\beta c + 1}}$$

In particular, omitting constant factors that depend on β , c , and L , we get

$$\epsilon_{\text{stab}} \lesssim \frac{T^{1-1/(\beta c + 1)}}{n}.$$

Proof. Let S and S' be two samples of size n differing in only a single example. Consider the gradient updates G_1, \dots, G_T and G'_1, \dots, G'_T induced by running SGM on sample S and S' , respectively. Let w_T and w'_T denote the corresponding outputs of SGM.

By Lemma 3.8, we have for every $t_0 \in \{1, \dots, n\}$,

$$\mathbb{E} |f(w_T; z) - f(w'_T; z)| \leq \frac{t_0}{n} + L \mathbb{E} [\delta_T \mid \delta_{t_0} = 0], \quad (3.7)$$

where $\delta_t = \|w_T - w'_T\|$. To simplify notation, let $\Delta_t = \mathbb{E} [\delta_T \mid \delta_{t_0} = 0]$. We will bound Δ_t a function of t_0 and then minimize for t_0 .

Toward this goal, observe that at step t , with probability $1 - 1/n$, the example selected by SGM is the same in both S and S' . In this case we have that $G_t = G'_t$ and we can use the $(1 + \alpha_t \beta)$ -expansivity of the update rule G_t which follows from our smoothness assumption via Lemma 3.7.1. With probability $1/n$ the selected example is different in which case we use that both G_t and G'_t are $\alpha_t L$ -bounded as a consequence of Lemma 3.3.

Hence, we can apply Lemma 2.5 and linearity of expectation to conclude that for every $t \geq t_0$,

$$\begin{aligned} \Delta_{t+1} &\leq \left(1 - \frac{1}{n}\right) (1 + \alpha_t \beta) \Delta_t + \frac{1}{n} \Delta_t + \frac{2\alpha_t L}{n} \\ &\leq \left(\frac{1}{n} + (1 - 1/n)(1 + c\beta/t)\right) \Delta_t + \frac{2cL}{tn} \\ &= \left(1 + (1 - 1/n)\frac{c\beta}{t}\right) \Delta_t + \frac{2cL}{tn} \\ &\leq \exp\left(\left(1 - 1/n\right)\frac{c\beta}{t}\right) \Delta_t + \frac{2cL}{tn}. \end{aligned}$$

Here we used that $1 + x \leq \exp(x)$ for all x .

Using the fact that $\Delta_{t_0} = 0$, we can unwind this recurrence relation from T down to $t_0 + 1$. This gives

$$\begin{aligned}
\Delta_T &\leq \sum_{t=t_0+1}^T \left\{ \prod_{k=t+1}^T \exp\left(\left(1 - \frac{1}{n}\right)\frac{\beta c}{k}\right) \right\} \frac{2cL}{tn} \\
&= \sum_{t=t_0+1}^T \exp\left(\left(1 - \frac{1}{n}\right)\beta c \sum_{k=t+1}^T \frac{1}{k}\right) \frac{2cL}{tn} \\
&\leq \sum_{t=t_0+1}^T \exp\left(\left(1 - \frac{1}{n}\right)\beta c \log\left(\frac{T}{t}\right)\right) \frac{2cL}{tn} \\
&= \frac{2cL}{n} T^{\beta c(1-1/n)} \sum_{t=t_0+1}^T t^{-\beta c(1-1/n)-1} \\
&\leq \frac{1}{(1-1/n)\beta c} \frac{2cL}{n} \left(\frac{T}{t_0}\right)^{\beta c(1-1/n)} \\
&\leq \frac{2L}{\beta(n-1)} \left(\frac{T}{t_0}\right)^{\beta c},
\end{aligned}$$

Plugging this bound into (3.7), we get

$$\mathbb{E} |f(w_T; z) - f(w'_T; z)| \leq \frac{t_0}{n} + \frac{2L^2}{\beta(n-1)} \left(\frac{T}{t_0}\right)^{\beta c}.$$

Letting $q = \beta c$, the right hand side is approximately minimized when

$$t_0 = (2cL^2)^{\frac{1}{q+1}} T^{\frac{q}{q+1}}.$$

This setting gives us

$$\mathbb{E} |f(w_T; z) - f(w'_T; z)| \leq \frac{1+1/q}{n-1} (2cL^2)^{\frac{1}{q+1}} T^{\frac{q}{q+1}} = \frac{1+1/\beta c}{n-1} (2cL^2)^{\frac{1}{\beta c+1}} T^{\frac{\beta c}{\beta c+1}}.$$

Since the bound we just derived holds for all S, S' and z , we immediately get the claimed upper bound on the uniform stability. \square

4 Stability-inducing operations

In light of our results, it makes sense to analyse for operations that increase the stability of the stochastic gradient method. We show in this section that pleasingly several popular heuristics and methods indeed improve the stability of SGM. Our rather straightforward analyses both to strengthen the bounds we previously obtained and provides an explanation of empirical success of the methods we overview.

Weight Decay and Regularization. Weight decay is a simple and effective that often improves generalization [20].

Definition 4.1. Let $f: \Omega \rightarrow \Omega$, be a differentiable function. We define the gradient update with weight decay at rate μ as $G_{f,\mu,\alpha}(w) = (1 - \alpha\mu)w - \alpha\nabla f(w)$.

It is easy to verify that the above update rule is equivalent to performing a gradient update on the ℓ_2 -regularized objective $g(w) = f(w) + \frac{\mu}{2}\|w\|^2$.

Lemma 4.2. Assume that f is β -smooth. Then, $G_{f,\mu,\alpha}$ is $(1 + \alpha(\beta - \mu))$ -expansive.

Proof. Let $G = G_{f,\mu,\alpha}$. By triangle inequality and our smoothness assumption,

$$\begin{aligned} \|G(v) - G(w)\| &\leq (1 - \alpha\mu)\|v - w\| + \alpha\|\nabla f(w) - \nabla f(v)\| \\ &\leq (1 - \alpha\mu)\|v - w\| + \alpha\beta\|w - v\| \\ &= (1 - \alpha\mu + \alpha\beta)\|v - w\|. \end{aligned}$$

□

The above lemma shows as that a regularization parameter μ counters a smoothness parameter β . Once $r > \beta$, the gradient update with decay becomes contractive. Any theorem we proved in previous sections that has a dependence on β leads to a corresponding theorem for stochastic gradient with weight decay in which β is replaced with $\beta - \mu$.

Gradient Clipping. It is common when training deep neural networks to enforce bounds on the norm of the gradients encountered by SGD. This is often done by either truncation, scaling, or dropping of examples that cause an exceptionally large value of the gradient norm. Any such heuristic directly leads to a bound on the Lipschitz parameter L that appears in our bounds. It is also easy to introduce a varying Lipschitz parameter L_t to account for possibly different values.

Dropout. Dropout [39] is a popular and effective heuristic for preventing large neural networks from overfitting. Here we prove that, indeed, dropout improves all of our stability bounds generically. From the point of view of stochastic gradient descent, dropout is equivalent to setting a fraction of the gradient weights to zero. That is, instead of updating with a stochastic gradient $\nabla f(w; z)$ we instead update with a perturbed gradient $D\nabla f(w; z)$ which is typically identical to $\nabla f(w; z)$ in some of the coordinates and equal to 0 on the remaining coordinates, although our definition is a fair bit more general.

Definition 4.3. We say that a randomized map $D: \Omega \rightarrow \Omega$ is a dropout operator with dropout rate s if for every $v \in D$ we have $\mathbb{E}\|Dv\| = s\|v\|$. For a differentiable function $f: \Omega \rightarrow \Omega$, we let $DG_{f,\alpha}$ denote the dropout gradient update defined as $DG_{f,\alpha}(v) = v - \alpha P\nabla f(v)$

As expected, the dropout update improves the effective Lipschitz constant of the objective function.

Lemma 4.4. Assume that f is L -Lipschitz. Then, the dropout update $DG_{f,\alpha}$ with dropout rate s is $(s\alpha L)$ -bounded.

Proof. By our Lipschitz assumption and linearity of expectation,

$$\mathbb{E} \|G_{f,\alpha}(v) - v\| = \alpha \mathbb{E} \|D\nabla f(v)\| = \alpha s \mathbb{E} \|\nabla f(v)\| \leq \alpha s L, .$$

□

From this lemma we can obtain various corollaries by replacing L with sL in our theorems.

Projections and Proximal Steps. Related to regularization, there are many popular updates which follow a stochastic gradient update with a projection onto a set or some statistical shrinkage operation. The vast majority of these operations can be understood as applying a proximal-point operation associated with a convex function. Similar to the gradient operation, we can define the proximal update rule.

Definition 4.5. For a nonnegative step size $\alpha \geq 0$ and a function $f: \Omega \rightarrow \mathbb{R}$, we define the proximal update rule $P_{f,\alpha}$ as

$$P_{f,\alpha}(w) = \arg \min_v \frac{1}{2} \|w - v\|^2 + \alpha f(v). \quad (4.1)$$

For example, Euclidean projection is the proximal point operation associated with the indicator of the associated set. Soft-thresholding is the proximal point operator associated with the ℓ_1 -norm. For more information, see the surveys by Combettes and Wajs [6] or Parikh and Boyd [32].

An elementary proof of the following Lemma, due to Rockafellar [35], can be found in the appendix.

Lemma 4.6. If f is convex, the proximal update (4.1) is 1-expansive.

In particular, this Lemma implies that the Euclidean projection onto a convex set is 1-expansive. Note that in many important cases, proximal operators are actually contractive. That is, they are η -expansive with $\eta < 1$. An notable example is when $f(\cdot)$ is the Euclidean norm for which the update rule is η -expansive with $\eta = (1 + \alpha)^{-1}$. So stability can be induced by the choice of an appropriate prox-operation, which can always be interpreted as some form of regularization.

Model Averaging. Model averaging refers to the idea of averaging out the iterates w_t obtained by a run of SGD. In convex optimization, model averaging is sometimes observed to lead to better empirical performance of SGM and closely replated updates such as the Perceptron [10]. Here we show that model averaging improves our bound for the convex optimization by a constant factor.

Theorem 4.7. Assume that $f: \Omega \rightarrow [0, 1]$ is a decomposable convex L -Lipschitz β -smooth function and that we run SGD with step sizes $\alpha_t \leq \alpha \leq 2/\beta$ for T steps. Then, the average of the first T iterates of SGD has uniform stability of $\epsilon_{\text{stab}} \leq \frac{\alpha T L^2}{n}$.

Proof. Let $\bar{w}_T = \frac{1}{T} \sum_{t=1}^T w_t$ denote the average of the stochastic gradient iterates. Since

$$w_t = \sum_{k=1}^t \alpha \nabla f(w_k; (x_k, y_k)),$$

we have

$$\bar{w}_T = \alpha \sum_{t=1}^T \frac{T-t+1}{T} \nabla f(w_k; (x_k, y_k))$$

Using Lemma 3.8, the deviation between \bar{w}_t and \bar{w}'_t obeys

$$\delta_t \leq (1 - 1/n)\delta_{t-1} + \frac{1}{n} \left(\delta_{t-1} + 2\alpha L \frac{T-t+1}{T} \right).$$

which implies

$$\delta_T \leq \frac{2\alpha L}{n} \sum_{t=1}^T \frac{T-t+1}{T} = \frac{\alpha L(T+1)}{n}.$$

Since f is L -Lipschitz, we have

$$\mathbb{E} |f(\bar{w}_T) - f(\bar{w}'_T)| \leq L \|\bar{w}_T - \bar{w}'_T\| \leq \frac{\alpha(T+1)L^2}{n}.$$

Here the expectation is taken over the algorithm and hence the claim follows by our definition of uniform stability. \square

5 Convex risk minimization

We now outline how our generalization bounds lead to bounds on the population risk achieved by SGM in the convex setting. We restrict our attention to the convex case where we can contrast against known results. The main feature of our results is that we show that one can achieve bounds comparable or perhaps better than known results on stochastic gradient for risk minimization by running for multiple passes over the data set.

The key to the analysis in this section is to decompose the risk estimates into an *optimization error* term and a stability term. The optimization error designates how closely we optimize the empirical risk or a proxy of the empirical risk. By optimizing with stochastic gradient, we will be able to balance this optimization accuracy against how well we generalize. These results are inspired by the work of Bousquet and Bottou who provided similar analyses for SGM based on uniform convergence [3]. However, our stability results will yield sharper bounds.

Throughout this section, our risk decomposition works as follows. We define the *optimization error* to be the gap between the empirical risk and minimum empirical risk in expectation:

$$\epsilon_{\text{opt}}(w) \stackrel{\text{def}}{=} \mathbb{E} [R_S[w] - R_S[w_\star^S]] \quad \text{where } w_\star^S = \arg \min_w R_S[w].$$

By Theorem 2.2, the expected risk of a w output by SGM is bounded as

$$\mathbb{E}[R[w]] \leq \mathbb{E}[R_S[w]] + \epsilon_{\text{stab}} \leq \mathbb{E}[R_S[w_\star^S]] + \epsilon_{\text{opt}}(w) + \epsilon_{\text{stab}}$$

In general, the optimization error decreases with the number of SGM iterations while the stability increases. Balancing these two terms will thus provide a reasonable excess risk against the empirical risk minimizer. Note that our analysis involves the expected minimum empirical risk which could be considerably smaller than the minimum risk. However, as we now show, it can never be larger.

Lemma 5.1. *Let w_\star denote the minimizer of the population risk and w_\star^S denote the minimizer of the empirical risk given a sampled data set S . Then $\mathbb{E}[R_S[w_\star^S]] \leq R[w_\star]$.*

Proof.

$$\begin{aligned} R[w_\star] &= \inf_w R[w] = \inf_w \mathbb{E}_z[f(w; z)] \\ &= \inf_w \mathbb{E}_S \left[\frac{1}{n} \sum_{i=1}^n f(w; z_i) \right] \\ &\geq \inf_w \mathbb{E}_S \left[\frac{1}{n} \sum_{i=1}^n f(w_\star^S; z_i) \right] \\ &= \mathbb{E}_S \left[\frac{1}{n} \sum_{i=1}^n f(w_\star^S; z_i) \right] = \mathbb{E}[R_S[w_\star^S]]. \end{aligned}$$

□

To analyze the optimization error, we will make use of a classical result due to Nemirovski and Yudin [28].

Theorem 5.2. *Assume we run stochastic gradient descent with constant stepsize α on a convex function*

$$R[w] = \mathbb{E}_z[f(w; z)].$$

Assume further that $\|\nabla f(w; z)\| \leq L$ and $\|w_0 - w_\star\| \leq D$ for some minimizer w_\star of J . Let \bar{w}_T denote the average of the T iterates of the algorithm. Then we have

$$R[\bar{w}_T] \leq R[w_\star] + \frac{1}{2} \frac{D^2}{T\alpha} + \frac{1}{2} L^2 \alpha.$$

The upper bound stated in the previous theorem is known to be tight even if the function is β -smooth [28]

If we plug in the population risk for J in the previous theorem, we directly obtain a generalization bound for SGM that holds when we make a single pass over the data. The theorem requires fresh samples from the distribution in each update step of SGM. Hence, given n data points, we cannot make more than n steps, and each sample must not be used more than once.

Corollary 5.3. *Let f be a convex loss function satisfying $\|\nabla f(w, z)\| \leq L$ and let w_\star be a minimizer of the population risk $R[w] = \mathbb{E}_z f(w; z)$. Suppose we make a single pass of SGM over the sample $S = (z_1, \dots, z_n)$ with a suitably chosen fixed step size starting from a point w_0 that satisfies $\|w_0 - w_\star\| \leq D$. Then, the average \bar{w}_n of the iterates satisfies*

$$\mathbb{E}[R[\bar{w}_n]] \leq R[w_\star] + \frac{DL}{\sqrt{n}}. \tag{5.1}$$

We now contrast this bound with what follows from our results.

Proposition 5.4. *Let $S = (z_1, \dots, z_n)$ be a sample of size n . Let f be a β -smooth convex loss function satisfying $\|\nabla f(w, z)\| \leq L$ and let w_\star^S be a minimizer of the empirical risk $R_S[w] =$*

$\frac{1}{n} \sum_{i=1}^n f(w; z_i)$. Suppose we run T steps of SGM with suitably chosen step size from a starting point w_0 that satisfies $\|w_0 - w_\star^S\| \leq D$. Then, the average \bar{w}_T over the iterates satisfies

$$\mathbb{E}[R[\bar{w}_T]] \leq \mathbb{E}[R_S[w_\star^S]] + \frac{DL}{\sqrt{n}} \sqrt{\frac{n+2T}{T}}.$$

Proof. On the one hand, applying Theorem 5.2 to the empirical risk R_S , we get

$$\epsilon_{\text{opt}}(\bar{w}_T) \leq \frac{1}{2} \frac{D^2}{T\alpha} + \frac{1}{2} L^2 \alpha.$$

Here, w_\star^S is an empirical risk minimizer. On the other hand, by our stability bound from Theorem 4.7,

$$\epsilon_{\text{stab}} \leq \frac{TL^2\alpha}{n}$$

Combining these two inequalities we have,

$$\mathbb{E}[R[\bar{w}_T]] \leq \mathbb{E}[R_{\text{emp}}[w_\star^S]] + \frac{1}{2} \frac{D^2}{T\alpha} + \frac{1}{2} L^2 \left(1 + \frac{2T}{n}\right) \alpha$$

Choosing α to be

$$\alpha = \frac{D\sqrt{n}}{L\sqrt{T(n+2T)}},$$

yields the bound provided in the proposition. \square

Note that the bound from our stability analysis is not directly comparable to Corollary 5.3 as we are comparing against the expected minimum empirical risk rather than the minimum risk. Lemma 5.1 implies that the excess risk in our bound is at most worse by a factor of $\sqrt{3}$ compared with Corollary 5.3 when $T = n$. Moreover, the excess risk in our bound tends to a factor merely $\sqrt{2}$ larger than the Nemirovski-Yudin bound as T goes to infinity. In contrast, the classical bound does not apply when $T > n$.

6 Future Work and Open Problems

Our analysis parts from much previous work in that we directly analyze the generalization performance of an algorithm rather than the solution of an optimization problem. In doing so we build on the toolkit usually used to prove that algorithms converge in objective value.

This approach could be more powerful than analyzing optimality conditions, as it may be easier to understand how each data point affects a procedure rather than an optimal solution. It also has the advantage that the generalization bound holds even if the algorithm fails to find a unique optimal solution as is common in non-convex problems.

In addition to this broader perspective on algorithms for learning, there are many exciting theoretical and empirical directions that we intend to pursue in future work. We close this manuscript by detailing some of the more concrete open problems and avenues for direct empirical validation.

High Probability Bounds The results in this paper are all *in expectation*. Similar to the well-known proofs of the stochastic gradient method, deriving bounds on the expected risk is relatively straightforward, but high probability bounds need more attention and care [26, 34]. In the case of stability, the standard techniques from Bousquet and Elisseeff require uniform stability on the order of $O(1/n)$ to apply exponential concentration inequalities like McDiarmid’s [4]. For larger values of the stability parameter ϵ_{stab} , it is more difficult to construct such high probability bounds. In our setting, things are further complicated by the fact that our algorithm is itself randomized, and thus a concentration inequality must be devised to account for both the randomness in the data and in the training algorithm. Since differential privacy and stability are closely related, one possibility is to derive concentration via an algorithmic method, similar to the one developed by Nissim and Stemmer [31].

Stability of the gradient method Since gradient descent can be considered a “limiting case” of the stochastic gradient descent method, one can use an argument like our Growth Recursion Lemma (Lemma 2.5) to analyze its stability. Such an argument provides an estimate of $\epsilon_{\text{stab}} \leq \frac{\alpha T}{n}$ where α is the step size and T is the number of iterations. Generic bounds for convex functions suggest that gradient descent achieves an optimization error of $O(1/T)$. Thus, a generalization bound of $O(1/\sqrt{n})$ is achievable, but at a computational complexity of $O(n^{1.5})$. SGM, on the other hand, achieves a generalization of $O(1/\sqrt{n})$ in time $O(n)$.

In the non-convex case, we are unable to prove any reasonable form of stability at all. We are unable to apply our “burn-in” argument to the gradient method, and thus trajectories may diverge exponentially quickly. Poor generalization behavior of gradient descent has been observed in practice, but lower bounds for this approach are necessary to rule out a stable implementation for non-convex machine learning.

Acceleration and momentum We have described how many of the best practices in neural net training can be understood as stability inducing operations. One very important technique that we did not discuss is *momentum*. In momentum methods, the update is a linear combination of the current iterate and the previous direction. For convex problems, momentum is known to decrease the number of iterations required by stochastic gradient descent [22]. For general nonlinear problems, is believed to decrease the number of iterations required to achieve low-training error [33, 37]. However, it is not clear that momentum adds stability. Indeed, in the case of convex optimization, momentum methods are less robust to noise than gradient methods [7, 23]. Thus, it is possible that momentum speeds up training but adversely impacts generalization.

Empirical Validation Most notably, we hope to investigate whether these theoretical developments parallel empirical behavior. For such an investigation, it is necessary to devise efficient methods to measure the stability exhibited by stochastic gradient. We can then determine if the models that generalize in practice are indeed the most stable ones. Moreover, we can verify whether minor algorithmic modifications that induce stability also result in the models which generalize best. We expect to perform such experiments in the near future.

Model Selection Another related avenue that bridges theory and practice is using stability as a method for model selection. In particular, our results imply that the models that train the fastest also generalize the best. This suggests that a heuristic for model selection would be to run many

different parameter settings and choose the model which results in the lowest training error most quickly. This idea is relatively simple to try in practice, and ideas from bandit optimization can be applied to efficiently search with this heuristic cost [15, 17]. From the theoretical perspective, understanding the sensibility of this heuristic would require understanding *lower bounds* for generalizability. Are there necessary conditions which state that models which take a long training time by SGM generalize less well than those with short training times?

High capacity models that train quickly If the models can be trained quickly via stochastic gradient, our results prove that these models will generalize. However, this manuscript provides no guidance as to how to build a model where training is stable *and* training error is low. Designing a family of models which both has high capacity and can be trained quickly would be of significant theoretical and practical interest.

Indeed, the capacity of models trained in current practice steadily increases as growing computational power makes it possible to effectively train larger models. It is not uncommon for some models, such as large neural networks, to have more free parameters than the size of the sample yet have rather small generalization error [19, 40]. In fact, sometimes increasing the model capacity even seems to *decrease* the generalization error [30]. Is it possible to understand this phenomena via stability? How can we find models which provably both have high capacity and train quickly?

Algorithm Design Finally, we note that stability may also provide new ideas for designing learning rules. There are a variety of successful methods in machine learning and signal processing that do not compute an exact stochastic gradient, yet are known to find quality stationary points in theory and practice [5]. Do the ideas developed in this paper provide new insights into how to design learning rules that accelerate the convergence and improve the generalization of SGM?

Acknowledgements

The authors would like to thank John Duchi, Vineet Gupta, Kevin Jamieson, Eric Price, Nati Srebro, and Oriol Vinyals for their insightful feedback and helpful suggestions.

References

- [1] S. Arora, R. Ge, T. Ma, and A. Moitra. Simple, efficient, and neural algorithms for sparse coding. In *Proceedings of the Conference on Learning Theory (COLT)*, 2015.
- [2] L. Bottou. Online algorithms and stochastic approximations. In D. Saad, editor, *Online Learning and Neural Networks*. Cambridge University Press, Cambridge, UK, 1998.
- [3] L. Bottou and O. Bousquet. The tradeoffs of large scale learning. In *Neural Information Processing Systems*, 2008.
- [4] O. Bousquet and A. Elisseeff. Stability and generalization. *Journal of Machine Learning Research*, 2:499–526, 2002.
- [5] J. Bucklew, T. G. Kurtz, and W. Sethares. Weak convergence and local stability properties of fixed step size recursive algorithms. *IEEE Transactions on Information Theory*, 39(3):966–978, 1993.

- [6] P. L. Combettes and V. R. Wajs. Signal recovery by proximal forward-backward splitting. *Multiscale Modeling and Simulation*, 4(4):1168–1200, 2005.
- [7] O. Devolder, F. Glineur, and Y. Nesterov. First-order methods of smooth convex optimization with inexact oracle. *Mathematical Programming*, 146(1-2):37–75, 2014.
- [8] L. P. Devroye and T. Wagner. Distribution-free performance bounds for potential function rules. *Information Theory, IEEE Transactions on*, 25(5):601–604, Sep 1979.
- [9] A. Elisseeff, T. Evgeniou, and M. Pontil. Stability of randomized learning algorithms. *Journal of Machine Learning Research*, 6:55–79, 2005.
- [10] Y. Freund and R. E. Schapire. Large margin classification using the perceptron algorithm. *Machine Learning*, 37(3):277–296, 1999.
- [11] R. Frostig, R. Ge, S. M. Kakade, and A. Sidford. Competing with the empirical risk minimizer in a single pass. In *Conference on Learning Theory (COLT)*, 2015.
- [12] S. Ghadimi and G. Lan. Stochastic first-and zeroth-order methods for nonconvex stochastic programming. *SIAM Journal on Optimization*, 23(4):2341–2368, 2013.
- [13] E. Hazan, A. Kalai, S. Kale, and A. Agarwal. Logarithmic regret algorithms for online convex optimization. In *Proceedings of the 19th Annual Conference on Learning Theory*, 2006.
- [14] E. Hazan and S. Kale. Beyond the regret minimization barrier: optimal algorithms for stochastic strongly-convex optimization. *Journal of Machine Learning Research*, 15(1):2489–2512, 2014.
- [15] K. Jamieson and A. Talwalkar. Non-stochastic best arm identification and hyperparameter optimization. Preprint available at [arXiv:1502.07943v1](https://arxiv.org/abs/1502.07943v1), 2015.
- [16] M. Janzamin, H. Sedghi, and A. Anandkumar. Generalization bounds for neural networks through tensor factorization. Preprint available at [arXiv:1506.08473](https://arxiv.org/abs/1506.08473), 2015.
- [17] Z. Karnin, T. Koren, and O. Somekh. Almost optimal exploration in multi-armed bandits. In *Proceedings of the 30th International Conference on Machine Learning (ICML-13)*, 2013.
- [18] M. J. Kearns and D. Ron. Algorithmic stability and sanity-check bounds for leave-one-out cross-validation. *Neural Computation*, 11(6):1427–1453, 1999.
- [19] A. Krizhevsky, I. Sutskever, and G. E. Hinton. Imagenet classification with deep convolutional neural networks. In *Proc. NIPS*, pages 1097–1105, 2012.
- [20] A. Krogh and J. A. Hertz. A simple weight decay can improve generalization. In *Proc. NIPS*, pages 950–957, 1992.
- [21] H. J. Kushner and G. G. Yin. *Stochastic Approximation and Recursive Algorithms and Applications*. Springer-Verlag, New York, second edition, 2003.
- [22] G. Lan. An optimal method for stochastic composite optimization. *Mathematical Programming*, 133(1-2):365–397, 2012.

- [23] L. Lessard, B. Recht, and A. Packard. Analysis and design of optimization algorithms via integral quadratic constraints. To appear in *SIAM Journal on Optimization*. Preprint available at [arxiv:1408.3595](https://arxiv.org/abs/1408.3595), 2014.
- [24] R. Livni, S. Shalev-Shwartz, and O. Shamir. On the computational efficiency of training neural networks. In *Advances in Neural Information Processing Systems*, pages 855–863, 2014.
- [25] S. Mukherjee, P. Niyogi, T. Poggio, and R. M. Rifkin. Learning theory: stability is sufficient for generalization and necessary and sufficient for consistency of empirical risk minimization. *Adv. Comput. Math.*, 25(1-3):161–193, 2006.
- [26] A. Nemirovski, A. Juditsky, G. Lan, and A. Shapiro. Robust stochastic approximation approach to stochastic programming. *SIAM Journal on Optimization*, 19(4):1574–1609, 2009.
- [27] A. Nemirovski and D. B. Yudin. On Cezari’s convergence of the steepest descent method for approximating saddle point of convex-concave functions. In *Soviet Mathematics Doklady*, volume 19, 1978.
- [28] A. Nemirovski and D. B. Yudin. *Problem complexity and method efficiency in optimization*. Wiley Interscience, 1983.
- [29] Y. Nesterov. *Introductory lectures on convex optimization*, volume 87 of *Applied Optimization*. Kluwer Academic Publishers, Boston, MA, 2004. A basic course.
- [30] B. Neyshabur, R. Tomioka, and N. Srebro. In search of the real inductive bias: On the role of implicit regularization in deep learning. In *In Proc. 3rd ICLR*, 2015.
- [31] K. Nissim and U. Stemmer. On the generalization properties of differential privacy. Preprint available at [arXiv:1504.05800](https://arxiv.org/abs/1504.05800), 2015.
- [32] N. Parikh and S. P. Boyd. Proximal algorithms. *Foundations and Trends in optimization*, 1(3):123–231, 2013.
- [33] B. T. Polyak. Introduction to optimization. *Optimization Software, Inc.*, 1987.
- [34] A. Rakhlin, O. Shamir, and K. Sridharan. Making gradient descent optimal for strongly convex stochastic optimization. In *Proceedings of the 29th International Conference on Machine Learning*, 2012. Extended version at [arxiv:1109.5647](https://arxiv.org/abs/1109.5647).
- [35] R. T. Rockafellar. Monotone operators and the proximal point algorithm. *SIAM Journal on Control and Optimization*, 14(5):877–898, 1976.
- [36] L. Rosasco and S. Villa. Learning with incremental iterative regularization. Preprint available at [arXiv:1405.0042v2](https://arxiv.org/abs/1405.0042v2), 2014.
- [37] D. E. Rumelhart, G. E. Hinton, and R. J. Williams. Learning internal representations by error propagation. In *Parallel Distributed Processing—Explorations in the Microstructure of Cognition*, pages 318–362. MIT Press, Cambridge, MA, 1986.
- [38] S. Shalev-Shwartz, O. Shamir, N. Srebro, and K. Sridharan. Learnability, stability and uniform convergence. *Journal of Machine Learning Research*, 11:2635–2670, 2010.

- [39] N. Srivastava, G. Hinton, A. Krizhevsky, I. Sutskever, and R. Salakhutdinov. Dropout: A simple way to prevent neural networks from overfitting. *The Journal of Machine Learning Research*, 15(1):1929–1958, 2014.
- [40] C. Szegedy, W. Liu, Y. Jia, P. Sermanet, S. Reed, D. Anguelov, D. Erhan, V. Vanhoucke, and A. Rabinovich. Going deeper with convolutions. In *Proceedings of the 2015 Conference on Computer Vision and Pattern Recognition*, 2015.

A Elementary properties of convex functions

Proof of Lemma 3.7.1. Let $G = G_{f,\alpha}$. By triangle inequality and our smoothness assumption,

$$\begin{aligned} \|G(v) - G(w)\| &\leq \|v - w\| + \alpha\|\nabla f(w) - \nabla f(v)\| \\ &\leq \|v - w\| + \alpha\beta\|w - v\| \\ &= (1 + \alpha\beta)\|v - w\|. \end{aligned} \quad \square$$

Proof of Lemma 3.7.2. Convexity and β -smoothness implies that the gradients are co-coercive, namely

$$\langle \nabla f(v) - \nabla f(w), v - w \rangle \geq \frac{1}{\beta} \|\nabla f(v) - \nabla f(w)\|^2. \quad (\text{A.1})$$

We conclude that

$$\begin{aligned} \|G_{f,\alpha}(v) - G_{f,\alpha}(w)\|^2 &= \|v - w\|^2 - 2\alpha\langle \nabla f(v) - \nabla f(w), v - w \rangle + \alpha^2\|\nabla f(v) - \nabla f(w)\|^2 \\ &\leq \|v - w\|^2 - \left(\frac{2\alpha}{\beta} - \alpha^2\right)\|\nabla f(v) - \nabla f(w)\|^2 \\ &\leq \|v - w\|^2. \end{aligned} \quad \square$$

Proof of Lemma 3.7.3. First, note that if f is γ strongly convex, then $\varphi(w) = f(w) - \frac{\gamma}{2}\|w\|^2$ is convex with $(\beta - \gamma)$ -smooth. Hence, applying (A.1) to φ yields the inequality

$$\langle \nabla f(v) - \nabla f(w), v - w, \rangle \geq \frac{\beta\gamma}{\beta + \gamma}\|v - w\|^2 + \frac{1}{\beta + \gamma}\|\nabla f(v) - \nabla f(w)\|^2$$

Using this inequality gives

$$\begin{aligned} \|G_{f,\alpha}(v) - G_{f,\alpha}(w)\| &= \|v - w\|^2 - 2\alpha\langle \nabla f(v) - \nabla f(w), v - w, \rangle + \alpha^2\|\nabla f(v) - \nabla f(w)\|^2 \\ &\leq \left(1 - 2\frac{\alpha\beta\gamma}{\beta + \gamma}\right)\|v - w\|^2 - \alpha\left(\frac{2}{\beta + \gamma} - \alpha\right)\|\nabla f(v) - \nabla f(w)\|^2. \end{aligned}$$

The lemma follows. □

Proof of Lemma 4.6. This proof is due to Rockafellar [35]. Define

$$P_\nu(w) = \arg \min_v \frac{1}{2\nu}\|w - v\|^2 + f(v). \quad (\text{A.2})$$

This is the proximal mapping associated with f . Define the map $Q_\nu(w) := w - P_\nu(w)$. Then, by the optimality conditions associated with (A.2), we have

$$\nu^{-1}Q_\nu(w) \in \partial f(P_\nu(w)).$$

By convexity of f , we then have

$$\langle P_\nu(v) - P_\nu(w), Q_\nu(v) - Q_\nu(w) \rangle \geq 0.$$

Using this inequality, we have

$$\begin{aligned} \|v - w\|^2 &= \|[P_\nu(v) - P_\nu(w)] + [Q_\nu(v) - Q_\nu(w)]\|^2 \\ &= \|P_\nu(v) - P_\nu(w)\|^2 + 2\langle P_\nu(v) - P_\nu(w), Q_\nu(v) - Q_\nu(w) \rangle + \|Q_\nu(v) - Q_\nu(w)\|^2 \\ &\geq \|P_\nu(v) - P_\nu(w)\|^2, \end{aligned}$$

thus completing the proof. □